

	Department Name Address	ALS	Paramedic
		Revision #	
		Implementation Date	
Protocol	2.2.4 Croup and Epiglottitis	Last Reviewed/Update Date	
Author / Owner		Medical Director	

Croup and epiglottitis are upper airway obstructions. Patient care should be focused on adequate oxygenation and ventilation during transport. Respiratory emergencies are life threatening in the pediatric population; it is critical to be alert for early signs of decompensation. Avoid agitating the child with suspected epiglottitis.

1. Baseline care standards.
2. Place patient in position of comfort (upright in parent's lap if possible).
3. Administer high flow oxygen by mask or blow by and monitor O₂ sats (may use humidified oxygen if available).
4. Monitor cardiac rhythm.
5. Focused history and assessment.
Differential Diagnosis:
 - Croup:
 - Viral infection usually in children 6 months to 4 years of age.
 - Mild fever - some hoarseness.
 - Barking "seal bark" cough.
 - Condition worsens at night.
 - Nasal flaring, tracheal tugging and intercostal retractions possible.
 - Restlessness.
 - Pale skin and cyanosis is possible.
 - Epiglottitis:
 - Bacterial infection usually in children 4 years of age and older.
 - Sudden onset of high fever.
 - Painful swallowing (child may be drooling due to difficult swallowing).
 - Child may sit in tripod position in attempt to open airway.
 - Nasal flaring, tracheal tugging, intercostal retractions, and stridor possible.
 - Child may appear to look very ill.
6. Administer:

- *Nebulized Racemic Epinephrine 0.25ml in 2.5cc Normal Saline* < **4yrs of age**
 - *Nebulized Racemic Epinephrine 0.5ml in 2.5cc Normal Saline* > **4yrs of age**
 - May consider Albuterol 1.25 mg in 3 ml of saline solution (0.625 mg in 3 ml of saline solution if < 30 kg)
7. Should the patient deteriorate be prepared to assist ventilations with intubation and BVM.
 8. Consult medical control for needle cricothyrotomy for difficult airway management.
 9. **Do not attempt to visualize internal airway in responsive patient.**
 10. **Establish IV Normal Saline TKO only in unresponsive patient.**

Medical Director's Signature

Date

Disclaimer:

The protocols have been developed by the North Dakota Department of Health are meant to be used as general guidance for developing protocols for individual emergency medical services agencies. These sample protocols are not meant to be medical or legal advice; nor do they establish standards of care. Each emergency medical services agency must tailor protocols based on their specific needs or capabilities. Local medical directors must be consulted with and approve any protocol(s) prior to becoming operational in an emergency medical services agency. directors must be consulted with and approve any protocol(s) prior to becoming operational in an emergency medical services agency. The North Dakota Department of Health make no representation on the accuracy of information contained herein and accepts no liability for any loss or damage arising from any content error or omission.